

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

## 1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Meloxidolor 5 mg/ml solution for injection for dogs, cats, cattle and pigs

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

One ml contains:

**Active substance:**

Meloxicam 5 mg

**Excipient:**

Ethanol 150 mg

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Solution for injection.

Clear yellow solution.

## 4. CLINICAL PARTICULARS

### 4.1 Target species

Dogs, cats, cattle (calves ) and pigs

### 4.2 Indications for use, specifying the target species

**Dogs:**

Alleviation of inflammation and pain in both acute and chronic musculo-skeletal disorders. Reduction of post-operative pain and inflammation following orthopaedic and soft tissue surgery.

**Cats:**

Reduction of post-operative pain after ovariohysterectomy and minor soft tissue surgery.

**Cattle:**

For use in acute respiratory infection with appropriate antibiotic therapy to reduce clinical signs in cattle.

For use in diarrhoea in combination with oral re-hydration therapy to reduce clinical signs in calves of over one week of age.

For the relief of post-operative pain following dehorning in calves.

**Pigs:**

For use in non-infectious locomotor disorders to reduce the symptoms of lameness and inflammation.

For the relief of post-operative pain associated with minor soft tissue surgery such as castration.

### 4.3 Contraindications

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

Do not use in dogs and cats suffering from gastrointestinal disorders such as irritation and haemorrhage, impaired hepatic, cardiac or renal function and haemorrhagic disorders.

Do not use in dogs and cats less than 6 weeks of age nor in cats of less than 2 kg.  
Do not use in cattle and pigs suffering from impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of ulcerogenic gastrointestinal lesions.  
For the treatment of diarrhoea in cattle, do not use in animals of less than one week of age.  
Do not use in pigs less than 2 days old.  
See also section 4.7.

#### **4.4 Special warnings for each target species**

Treatment of piglets with Meloxidolor before castration reduces post-operative pain.

To obtain pain relief for cattle and pigs during surgery co-medication with an appropriate anaesthetic/sedative/analgesic is needed.

To obtain the best possible pain relieving effect for pigs post-surgery Meloxidolor should be administered 30 minutes before surgical intervention.  
Treatment of calves with Meloxidolor 20 minutes before dehorning reduces post-operative pain.  
Meloxidolor alone will not provide adequate pain relief during the dehorning procedure.

#### **4.5 Special precautions for use**

##### Special precautions for use in animals

Avoid use in any dehydrated, hypovolaemic or hypotensive animal, as there is a potential risk of renal toxicity. During anaesthesia, monitoring and fluid therapy should be considered as standard practice. Any oral follow-up therapy using meloxicam or other Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) should not be administered in cats, as appropriate dosage regimens for such follow-up treatments have not been established.

##### Special precautions to be taken by the person administering the veterinary medicinal product to animals

Accidental self-injection may give rise to pain. People with known hypersensitivity to NSAIDs should avoid contact with the veterinary medicinal product.

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

The veterinary medicinal product should not be administered by pregnant women or women of child-bearing potential as Meloxicam may be harmful to the foetus and unborn child.

#### **4.6 Adverse reactions (frequency and seriousness)**

##### For dogs and cats:

Typical adverse reactions of NSAIDs such as loss of appetite, vomiting, diarrhoea, faecal occult blood, lethargy and renal failure have rarely been reported. In very rare cases elevated liver enzymes have been reported.

In very rare cases, haemorrhagic diarrhoea, haematemesis and gastrointestinal ulceration have been reported. These adverse reactions occur generally within the first treatment week and are in most cases transient and disappear following termination of the treatment but in very rare cases may be serious or fatal.

In very rare cases anaphylactoid reactions may occur and should be treated symptomatically.

If adverse reactions occur, treatment should be discontinued and the advice of a veterinarian should be sought.

#### For cattle and pigs:

Only a slight transient swelling at the injection site following subcutaneous administration was observed in less than 10 % of the cattle treated in clinical studies.

In very rare cases anaphylactic reactions, which may be serious (including fatal), may occur and should be treated symptomatically.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

#### **4.7 Use during pregnancy, lactation or lay**

##### **Dogs and cats:**

Do not use in pregnant or lactating dogs or cats.

**Cattle:** Can be used during pregnancy.

**Pigs:** Can be used during pregnancy and lactation.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

##### For dogs and cats:

Other NSAIDs, diuretics, anticoagulants, aminoglycoside antibiotics and substances with high protein binding may compete for binding and thus lead to toxic effects. Meloxicam must not be administered in conjunction with other NSAIDs or glucocorticosteroids. Concurrent administration of potential nephrotoxic veterinary medicinal products should be avoided. In animals at anaesthetic risk (e.g. aged animals) intravenous or subcutaneous fluid therapy during anaesthesia should be taken into consideration. When anaesthesia and NSAID are concomitantly administered, a risk for renal function cannot be excluded.

Pre-treatment with anti-inflammatory substances may result in additional or increased adverse effects and accordingly a treatment-free period with such veterinary medicinal products should be observed for at least 24 hours before commencement of treatment. The treatment-free period, however, should take into account the pharmacological properties of the products used previously.

##### For cattle and pigs:

Do not administer concurrently with glucocorticosteroids, other non-steroidal anti-inflammatory drugs or with anticoagulant agents.

#### **4.9 Amounts to be administered and administration route**

##### Dogs:

Musculo-skeletal disorders:

Single subcutaneous injection at a dosage of 0.2 mg meloxicam/kg body weight (i.e. 0.4 ml/10 kg body weight).

Oral suspensions of meloxicam for dogs may be used for continuation of treatment at a dosage of 0.1 mg meloxicam/kg body weight, 24 hours after administration of the injection.

Reduction of post-operative pain (over a period of 24 hours):

Single intravenous or subcutaneous injection at a dosage of 0.2 mg meloxicam/kg body weight (i.e. 0.4 ml/10 kg body weight) before surgery, for example at the time of induction of anaesthesia.

#### Cats:

Reduction of post-operative pain:

Single subcutaneous injection at a dosage of 0.3 mg meloxicam/kg body weight (i.e. 0.06 ml/kg body weight) before surgery, for example at the time of induction of anaesthesia.

#### Cattle:

Single subcutaneous or intravenous injection at a dosage of 0.5 mg meloxicam/kg body weight (i.e. 10 ml/100 kg body weight) in combination with antibiotic therapy or with oral re-hydration therapy, as appropriate.

#### Pigs:

##### Locomotor disorders:

Single intramuscular injection at a dosage of 0.4 mg meloxicam/kg body weight (i.e. 2 ml/25 kg body weight). If required, a second administration of meloxicam can be given after 24 hours.

##### Reduction of post-operative pain:

Single intramuscular injection at a dosage of 0.4 mg meloxicam/kg body weight (i.e. 0.4 ml/5 kg body weight) before surgery.

Particular care should be taken with regard to the accuracy of dosing including the use of an appropriate dosing device and careful estimation of body weight.

Avoid introduction of contamination during use. The stopper should not be punctured more than 20 times.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

In case of overdose symptomatic treatment should be initiated.

#### **4.11 Withdrawal period(s)**

**Cattle:** Meat and offal: 15 days

**Pigs:** Meat and offal: 5 days

### **5. PHARMACOLOGICAL PROPERTIES**

Pharmacotherapeutic group: Antiinflammatory and antirheumatic products, non-steroids (oxicams)  
ATCvet code: QM01AC06

#### **5.1 Pharmacodynamic properties**

Meloxicam is a Non-Steroidal Anti-Inflammatory Drug (NSAID) of the oxicam class which acts by inhibition of prostaglandin synthesis, thereby exerting anti-inflammatory, anti-exudative, analgesic and antipyretic effects. It reduces leukocyte infiltration into the inflamed tissue. To a minor extent it also inhibits collagen-induced thrombocyte aggregation. *In vitro* and *in vivo* studies demonstrated that meloxicam inhibits cyclooxygenase-2 (COX-2) to a greater extent than cyclooxygenase-1 (COX-1). Meloxicam also has anti-endotoxic properties because it has been shown to inhibit production of thromboxane B<sub>2</sub> induced by *E. coli* endotoxin administration in calves and pigs.

#### **5.2 Pharmacokinetic particulars**

##### Absorption

Following subcutaneous administration, meloxicam is completely bioavailable and maximal mean plasma concentrations of 0.73 µg/ml in dogs and 1.1 µg/ml in cats were reached approximately 2.5 hours and 1.5 hours post administration, respectively.

After a single subcutaneous dose of 0.5 mg meloxicam/kg, C<sub>max</sub> values of 2.1 µg/ml were reached after

7.7 hours in young cattle.

Following single intramuscular doses of 0.4 mg meloxicam/kg, a  $C_{max}$  value of 1.1 to 1.5  $\mu\text{g/ml}$  was reached within 1 hour in pigs.

### Distribution

There is a linear relationship between the dose administered and plasma concentration observed in the therapeutic dose range in dogs and cats. More than 97 % of meloxicam is bound to plasma proteins.

The volume of distribution is 0.3 l/kg in dogs and 0.09 l/kg in cats.

In cattle and pigs, the highest meloxicam concentrations are to be found in liver and kidney.

Comparatively low concentrations are detectable in skeletal muscle and fat.

### Metabolism

Meloxicam is predominantly found in plasma. For dogs, cats and cattle it is also a major biliary excretion product whereas urine contains only traces of the parent compound.

In cattle, meloxicam is also a major excretion product in milk. In pigs, bile and urine contain only traces of the parent compound.

Five major metabolites were detected all having been shown to be pharmacologically inactive.

Meloxicam is metabolised to an alcohol, an acid derivative and to several polar metabolites. The main pathway of meloxicam biotransformation is oxidation.

### Elimination

In dogs and cats, Meloxicam is eliminated with a half-life of 24 hours. Approximately 75 % of the administered dose is eliminated via faeces and the remainder via urine in dogs.

In cats, the detection of metabolites from the parent compound in urine and faeces, but not in plasma is indicative for their rapid excretion. 21 % of the recovered dose is eliminated in urine (2 % as unchanged meloxicam, 19 % as metabolites) and 79 % in the faeces (49 % as unchanged meloxicam, 30 % as metabolites).

Meloxicam is eliminated with a half-life of 26 hours after subcutaneous injection in young cattle. In pigs, after intramuscular administration, the mean plasma elimination half-life is approximately 2.5 hours. Approximately 50 % of the administered dose is eliminated via urine and the remainder via faeces.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Ethanol

Poloxamer 188

Sodium chloride

Glycine

Sodium hydroxide

Hydrochloric acid

Glycofurol

Meglumine

Water for injections

### **6.2 Major incompatibilities**

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

### **6.3 Shelf life**

Shelf-life of the veterinary medicinal product as packaged for sale: 2 years.

Shelf-life after first opening the immediate packaging: 28 days

#### **6.4. Special precautions for storage**

This veterinary medicinal product does not require any special storage conditions.

#### **6.5 Nature and composition of immediate packaging**

Colourless type I glass vials of 10 ml, 20 ml or 100 ml, closed with a rubber stopper and sealed with an aluminium cap.

Multi-packs of 5 x 20 ml and 10 x 20 ml.

Not all pack sizes may be marketed.

#### **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

### **7. MARKETING AUTHORISATION HOLDER**

Le Vet Beheer B.V.  
Wilgenweg 7  
3421 TV Oudewater  
The Netherlands  
Tel: +31 (0)348 565858  
Fax: +31 (0)348 565454  
E-mail: info@levetpharma.com

### **8. MARKETING AUTHORISATION NUMBERS**

EU/2/13/148/001  
EU/2/13/148/002  
EU/2/13/148/003  
EU/2/13/148/008  
EU/2/13/148/009

### **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 22/04/2013

Date of last renewal: 15/03/2018

### **10. DATE OF REVISION OF THE TEXT**

Detailed information on this veterinary medicinal product is available on the website of the European Medicines Agency (<http://www.ema.europa.eu/>).

### **PROHIBITION OF SALE, SUPPLY AND/OR USE**

Not applicable.

## 1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Meloxidolor 20 mg/ml solution for injection for cattle, pigs and horses

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

One ml contains:

**Active substance:**

Meloxicam 20 mg

**Excipient:**

Ethanol 150 mg

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Solution for injection.

Clear yellow solution.

## 4. CLINICAL PARTICULARS

### 4.1 Target species

Cattle, pigs and horses

### 4.2 Indications for use, specifying the target species

**Cattle:**

For use in acute respiratory infection with appropriate antibiotic therapy to reduce clinical signs in cattle.

For use in diarrhoea in combination with oral re-hydration therapy to reduce clinical signs in calves of over one week of age and young, non-lactating cattle.

For adjunctive therapy in the treatment of acute mastitis, in combination with antibiotic therapy.

For the relief of post-operative pain following dehorning in calves.

**Pigs:**

For use in non-infectious locomotor disorders to reduce the symptoms of lameness and inflammation.

For adjunctive therapy in the treatment of puerperal septicaemia and toxæmia (mastitis-metritis-agalactia syndrome) with appropriate antibiotic therapy.

**Horses:**

For use in the alleviation of inflammation and relief of pain in both acute and chronic musculo-skeletal disorders.

For the relief of pain associated with equine colic.

### 4.3 Contraindications

See also section 4.7.

Do not use in horses less than 6 weeks of age.

Do not use in animals suffering from impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of ulcerogenic gastrointestinal lesions.



Do not use in cases of hypersensitivity to the active substance or to any of the excipients.  
For the treatment of diarrhoea in cattle, do not use in animals of less than one week of age

#### **4.4 Special warnings for each target species**

Treatment of calves with Meloxidolor 20 minutes before dehorning reduces post-operative pain. Meloxidolor alone will not provide adequate pain relief during the dehorning procedure. To obtain adequate pain relief during surgery co-medication with an appropriate analgesic is needed.

#### **4.5 Special precautions for use**

##### Special precautions for use in animals

If adverse reactions occur, treatment should be discontinued and the advice of a veterinarian should be sought.

Avoid use in very severely dehydrated, hypovolaemic or hypotensive animals which require parenteral rehydration, as there may be a potential risk of renal toxicity.

In case of inadequate relief of pain when used in the treatment of equine colic, careful re-evaluation of the diagnosis should be made as this could indicate the need for surgical intervention.

##### Special precautions to be taken by the person administering the veterinary medicinal product to animals

Accidental self-injection may give rise to pain. People with known hypersensitivity to Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) should avoid contact with the veterinary medicinal product.

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

The veterinary medicinal product should not be administered by pregnant women or women of child-bearing potential as Meloxicam may be harmful to the foetus and unborn child.

#### **4.6 Adverse reactions (frequency and seriousness)**

In cattle and pigs, subcutaneous, intramuscular as well as intravenous administration is well tolerated; only a slight transient swelling at the injection site following subcutaneous administration was observed in less than 10 % of the cattle treated in clinical studies.

In horses, in rare cases a transient swelling at the injection site can occur but resolves without intervention.

In very rare cases anaphylactoid reactions, which may be serious (including fatal), may occur and should be treated symptomatically.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reactions)
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

#### **4.7 Use during pregnancy, lactation or lay**

**Cattle and pigs:** Can be used during pregnancy and lactation.

**Horses:** Do not use in pregnant or lactating mares.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

Do not administer concurrently with glucocorticosteroids, other non-steroidal anti-inflammatory drugs or with anticoagulant agents.

#### 4.9 Amounts to be administered and administration route

##### **Cattle:**

Single subcutaneous or intravenous injection at a dosage of 0.5 mg meloxicam/kg body weight (i.e. 2.5 ml/100 kg body weight) in combination with antibiotic therapy or with oral re-hydration therapy, as appropriate.

##### **Pigs:**

Single intramuscular injection at a dosage of 0.4 mg meloxicam/kg body weight (i.e. 2 ml/100 kg body weight) in combination with antibiotic therapy, as appropriate. If required, a second administration of meloxicam can be given after 24 hours.

##### **Horses:**

Single intravenous injection at a dosage of 0.6 mg meloxicam/kg body weight (i.e. 3 ml/100 kg body weight).

Avoid introduction of contamination during use. The stopper should not be punctured more than 20 times.

#### 4.10 Overdose (symptoms, emergency procedures, antidotes if necessary)

In case of overdose symptomatic treatment should be initiated.

#### 4.11 Withdrawal period(s)

**Cattle:** Meat and offal: 15 days;  
Milk: 5 days

**Pigs:** Meat and offal: 5 days

**Horses:** Meat and offal: 5 days.

Not authorised for use in horses producing milk for human consumption.

### 5. PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group: Antiinflammatory and antirheumatic products, non-steroids (oxicams).  
ATCvet code: QM01AC06

#### 5.1 Pharmacodynamic properties

Meloxicam is a Non-Steroidal Anti-Inflammatory Drug (NSAID) of the oxicam class which acts by inhibition of prostaglandin synthesis, thereby exerting anti-inflammatory, anti-exudative, analgesic and antipyretic effects. It reduces leukocyte infiltration into the inflamed tissue. To a minor extent it also inhibits collagen-induced thrombocyte aggregation. Meloxicam also has anti-endotoxic properties because it has been shown to inhibit production of thromboxane B<sub>2</sub> induced by *E. coli* endotoxin administration in calves, lactating cows and pigs.

#### 5.2 Pharmacokinetic particulars

##### Absorption

After a single subcutaneous dose of 0.5 mg meloxicam/kg, C<sub>max</sub> values of 2.1 µg/ml and 2.7 µg/ml were reached after 7.7 hours and 4 hours in young cattle and lactating cows, respectively.

After two intramuscular doses of 0.4 mg meloxicam/kg, a C<sub>max</sub> value of 1.9 µg/ml was reached after 1 hour in pigs.

### Distribution

More than 98 % of meloxicam is bound to plasma proteins. The highest meloxicam concentrations are to be found in liver and kidney. Comparatively low concentrations are detectable in skeletal muscle and fat.

### Metabolism

Meloxicam is predominantly found in plasma. In cattle, meloxicam is also a major excretion product in milk and bile whereas urine contains only traces of the parent compound. In pigs, bile and urine contain only traces of the parent compound. Meloxicam is metabolised to an alcohol, an acid derivative and to several polar metabolites. All major metabolites have been shown to be pharmacologically inactive. The metabolism in horses has not been investigated.

### Elimination

Meloxicam is eliminated with a half-life of 26 hours and 17.5 hours after subcutaneous injection in young cattle and lactating cows, respectively.

In pigs, after intramuscular administration the mean plasma elimination half-life is approximately 2.5 hours.

In horses, after intravenous injection meloxicam is eliminated with a terminal half-life of 8.5 hours. Approximately 50 % of the administered dose is eliminated via urine and the remainder via faeces.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Ethanol  
Poloxamer 188  
Macrogol 300  
Glycine  
Disodium edetate  
Sodium hydroxide  
Hydrochloric acid  
Meglumine  
Water for injections

### **6.2 Major incompatibilities**

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

### **6.3 Shelf life**

Shelf-life of the veterinary medicinal product as packaged for sale: 2 years.

Shelf-life after first opening the immediate packaging: 28 days

### **6.4. Special precautions for storage**

This veterinary medicinal product does not require any special storage conditions.

### **6.5 Nature and composition of immediate packaging**

Colourless type I glass vial containing 50 ml or 100 ml, closed with a rubber stopper and sealed with an aluminium cap.

Multi-pack of 12 x 100ml

Not all pack sizes may be marketed.

**6.6 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

**7. MARKETING AUTHORISATION HOLDER**

Le Vet Beheer B.V. Wilgenweg 7  
3421 TV Oudewater  
The Netherlands  
tel: +31 (0)348 565858  
fax: +31 (0)348 565454  
e-mail: info@levetpharma.com

**8. MARKETING AUTHORISATION NUMBER(S)**

EU/2/13/148/004  
EU/2/13/148/005  
EU/2/13/148/010

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 22/04/2013  
Date of last renewal: 15/03/2018

**10. DATE OF REVISION OF THE TEXT**

Detailed information on this veterinary medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu/>.

**PROHIBITION OF SALE, SUPPLY AND/OR USE**

Not applicable.

**ANNEX II**

- A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. STATEMENT OF THE MRLs**

## A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE

### Name and address of the manufacturer responsible for batch release

Produlab Pharma B.V.  
Forellenweg 16  
4941 SJ Raamsdonksveer  
The Netherlands

## B. CONDITIONS OR RESTRICTIONS OF THE MARKETING AUTHORISATION REGARDING SUPPLY AND USE

Veterinary medicinal product subject to prescription.

## C. STATEMENT OF THE MRLs

The active substance in Meloxidolor is an allowed substance as described in table 1 of the annex to Commission Regulation (EU) No 37/2010:

Pharmacologically active substance	Marker residue	Animal species	MRL	Target tissues	Other provisions	Therapeutic classification
Meloxicam	Meloxicam	Bovine, caprine, porcine, rabbit, Equidae	20 µg/kg 65 µg/kg 65 µg/kg	Muscle Liver Kidney	NO ENTRY	Anti-inflammatory agents/Non-steroidal anti-inflammatory agents
		Bovine, caprine	15 µg/kg	Milk		

The excipients listed in section 6.1 of the SPC are either allowed substances for which table 1 of the annex to Commission Regulation (EU) No 37/2010 indicates that no MRLs are required or are considered as not falling within the scope of Regulation (EC) No 470/2009 when used as in this veterinary medicinal product.

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**



**PARTICULARS TO APPEAR ON THE OUTER PACKAGE AND THE IMMEDIATE PACKAGE**

**Carton for the 10 ml, 20 ml and 100 ml**

**Label for 100 ml**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Meloxidolor 5 mg/ml solution for injection for dogs, cats, cattle and pigs  
meloxicam

**2. STATEMENT OF ACTIVE SUBSTANCES**

Meloxicam 5 mg/ml

**3. PHARMACEUTICAL FORM**

Solution for injection

**4. PACKAGE SIZE**

10 ml  
20 ml  
100 ml

**5. TARGET SPECIES**

Dogs, cats, cattle (calves) and pigs

**6. INDICATION(S)**

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Dogs, Cattle: Single subcutaneous or intravenous injection  
Cats: Single subcutaneous injection  
Pigs: Single intramuscular injection  
Read the package leaflet before use

**8. WITHDRAWAL PERIOD(S)**

Withdrawal period:

**Cattle:** Meat and offal: 15 days

**Pigs:** Meat and offal: 5 days

**9. SPECIAL WARNING(S), IF NECESSARY**

Read the package leaflet before use.

**10. EXPIRY DATE**

EXP (month/year)  
Once broached, use by....

**11. SPECIAL STORAGE CONDITIONS**

Read the package leaflet before use.

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: read package leaflet.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only. To be supplied only on veterinary prescription.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Le Vet Beheer B.V.  
Wilgenweg 7  
3421 TV Oudewater  
The Netherlands

**16. MARKETING AUTHORISATION NUMBER(S)**

EU/2/13/148/001  
EU/2/13/148/002  
EU/2/13/148/003  
EU/2/13/148/008  
EU/2/13/148/009

**17. MANUFACTURER’S BATCH NUMBER**

Batch {number}

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**Label for 10 ml and 20 ml**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Meloxidolor 5 mg/ml solution for injection for dogs, cats, cattle and pigs  
meloxicam

**2. QUANTITY OF THE ACTIVE SUBSTANCE(S)**

Meloxicam 5 mg/ml

**3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES**

10 ml  
20 ml

**4. ROUTE(S) OF ADMINISTRATION**

Cattle: SC, IV  
Pigs: IM  
Dogs: IV or SC  
Cats: SC

**5. WITHDRAWAL PERIOD(S)**

Withdrawal period:  
**Cattle:** Meat and offal: 15 days  
**Pigs:** Meat and offal: 5 days

**6. BATCH NUMBER**

Batch {number}

**7. EXPIRY DATE**

EXP (month/year)  
Once broached, use by ....

**8. THE WORDS "FOR ANIMAL TREATMENT ONLY"**

For animal treatment only.

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

**Carton for 50 ml and 100 ml**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Meloxidolor 20 mg/ml solution for injection for cattle, pigs and horses  
meloxicam

**2. STATEMENT OF ACTIVE SUBSTANCES**

Meloxicam 20 mg/ml

**3. PHARMACEUTICAL FORM**

Solution for injection

**4. PACKAGE SIZE(S)**

50 ml  
100 ml

**5. TARGET SPECIES**

Cattle, pigs and horses

**6. INDICATION(S)**

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Cattle: single subcutaneous or intravenous injection  
Pigs: single intramuscular injection  
Horses: single intravenous injection  
Read the package leaflet before use.

**8. WITHDRAWAL PERIOD(S)**

Withdrawal period:  
**Cattle:** meat and offal: 15 days; milk: 5 days  
**Pigs, horses:** meat and offal: 5 days  
Not authorised for use in horses producing milk for human consumption.

**9. SPECIAL WARNING(S), IF NECESSARY**

Read the package leaflet before use.

**10. EXPIRY DATE**

EXP {month/year}  
Once broached, use by ....

**11. SPECIAL STORAGE CONDITIONS**

Read the package leaflet before use.

**12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: read package leaflet.

**13. THE WORDS "FOR ANIMAL TREATMENT ONLY" AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, if applicable**

For animal treatment only. To be supplied only on veterinary prescription.

**14. THE WORDS "KEEP OUT OF SIGHT AND REACH OF CHILDREN"**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Le Vet Beheer B.V.  
Wilgenweg 7  
3421 TV Oudewater  
The Netherlands

**16. MARKETING AUTHORISATION NUMBER(S)**

EU/2/13/148/004  
EU/2/13/148/005  
EU/2/13/148/010

**17. MANUFACTURER'S BATCH NUMBER**

Batch {number}

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE**

{Label for 100 ml}

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Meloxidolor 20 mg/ml solution for injection for cattle, pigs and horses  
meloxicam

**2. STATEMENT OF ACTIVE SUBSTANCES**

Meloxicam 20 mg/ml

**3. PHARMACEUTICAL FORM**

Solution for injection

**4. PACKAGE SIZES**

100 ml

**5. TARGET SPECIES**

Cattle, pigs and horses

**6. INDICATION(S)**

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

**8. WITHDRAWAL PERIOD(S)**

Withdrawal period:

Cattle: meat and offal: 15 days; milk: 5 days

Pigs, horses: meat and offal: 5 days

Not authorised for use in horses producing milk for human consumption.

**9. SPECIAL WARNING(S), IF NECESSARY**

Read the package leaflet before use.

**10. EXPIRY DATE**

EXP {month/year}

Once broached, use by

**11. SPECIAL STORAGE CONDITIONS**

Read the package leaflet before use.

**12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: read package leaflet.

**13. THE WORDS "FOR ANIMAL TREATMENT ONLY" AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, if applicable**

For animal treatment only. To be supplied only on veterinary prescription.

**14. THE WORDS "KEEP OUT OF SIGHT AND REACH OF CHILDREN"**

Keep out of the sight and reach of children

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Le Vet Beheer B.V.  
Wilgenweg 7  
3421 TV Oudewater  
The Netherlands

**16. MARKETING AUTHORISATION NUMBERS**

EU/2/13/148/005  
EU/2/13/148/010

**17. MANUFACTURER'S BATCH NUMBER**

Batch {number}

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Meloxidolor 20 mg/ml solution for injection for cattle, pigs and horses  
meloxicam

**2. QUANTITY OF THE ACTIVE SUBSTANCE(S)**

Meloxicam 20 mg/ml

**3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES**

50 ml

**4. ROUTE(S) OF ADMINISTRATION**

**Cattle:** SC or IV  
**Pigs:** IM  
**Horses:** IV

**5. WITHDRAWAL PERIOD(S)**

Withdrawal period:

Cattle: meat and offal: 15 days; milk: 5 days

Pigs, horses: meat and offal: 5 days

Not authorised for use in horses producing milk for human consumption.

**6. BATCH NUMBER**

Batch {number}

**7. EXPIRY DATE**

EXP {month/year}

Once broached, use by...

**8. THE WORDS "FOR ANIMAL TREATMENT ONLY"**

For animal treatment only.



**B. PACKAGE LEAFLET**

## PACKAGE LEAFLET:

### Meloxidolor 5 mg/ml solution for injection for dogs, cats, cattle and pigs

#### **1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder:

Le Vet Beheer B.V.  
Wilgenweg 7  
3421 TV Oudewater  
The Netherlands

Manufacturer responsible for the batch release:

Produlab Pharma B.V.  
Forellenweg 16  
4941 SJ Raamsdonksveer  
The Netherlands

#### **2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Meloxidolor 5 mg/ml solution for injection for dogs, cats, cattle and pigs  
meloxicam

#### **3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENTS**

One ml contains:

**Active substance:**

Meloxicam 5 mg

**Excipient:**

Ethanol 150 mg

Clear yellow solution.

#### **4. INDICATION(S)**

Dogs:

Alleviation of inflammation and pain in both acute and chronic musculo-skeletal disorders. Reduction of post-operative pain and inflammation following orthopaedic and soft tissue surgery.

Cats:

Reduction of post-operative pain after ovariohysterectomy and minor soft tissue surgery.

Cattle:

For use in acute respiratory infection with appropriate antibiotic therapy to reduce clinical signs in cattle.

For use in diarrhoea in combination with oral re-hydration therapy to reduce clinical signs in calves of over one week of age.

For the relief of post-operative pain following dehorning in calves.

Pigs:

For use in non-infectious locomotor disorders to reduce the symptoms of lameness and inflammation.  
For the relief of post-operative pain associated with minor soft tissue surgery such as castration.

## **5. CONTRAINDICATIONS**

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

Do not use in dogs and cats suffering from gastrointestinal disorders such as irritation and haemorrhage, impaired hepatic, cardiac or renal function and haemorrhagic disorders.  
Do not use in dog and cats less than 6 weeks of age nor in cats of less than 2 kg.

Do not use in cattle and pigs suffering from impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of ulcerogenic gastrointestinal lesions.  
For the treatment of diarrhoea in cattle, do not use in animals of less than one week of age.  
Do not use in pigs less than 2 days old.  
See also Section 12.

## **6. ADVERSE REACTIONS**

For dogs and cats:

Typical adverse reactions of Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) such as loss of appetite, vomiting, diarrhoea, faecal occult blood, lethargy and renal failure have rarely been reported. In very rare cases elevated liver enzymes have been reported.

In very rare cases, haemorrhagic diarrhoea, haematemesis and gastrointestinal ulceration have been reported. These side effects occur generally within the first treatment week and are in most cases transient and disappear following termination of the treatment but in very rare cases may be serious or fatal.

In very rare cases anaphylactoid reactions may occur and should be treated symptomatically.

For cattle and pigs:

Only a slight transient swelling at the injection site following subcutaneous administration was observed in less than 10 % of the cattle treated in clinical studies.

In very rare cases anaphylactic reactions, which may be serious (including fatal), may occur and should be treated symptomatically.

If adverse reactions occur, treatment should be discontinued and the advice of a veterinarian should be sought.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reactions)
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

## **7. TARGET SPECIES**

Dogs, cats, cattle (calves) and pigs

## **8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

### **Dosage for each species**

#### **Dogs:**

Musculo-skeletal disorders:

Single subcutaneous injection at a dosage of 0.2 mg meloxicam/kg body weight (i.e. 0.4 ml/10 kg body weight).

Oral suspensions of meloxicam for dogs may be used for continuation of treatment at a dosage of 0.1 mg meloxicam/kg body weight, 24 hours after administration of the injection.

Reduction of post-operative pain (over a period of 24 hours):

Single intravenous or subcutaneous injection at a dosage of 0.2 mg meloxicam/kg body weight (i.e. 0.4 ml/10 kg body weight) before surgery, for example at the time of induction of anaesthesia.

#### **Cats:**

Reduction of post-operative pain:

Single subcutaneous injection at a dosage of 0.3 mg meloxicam/kg body weight (i.e. 0.06 ml/kg body weight) before surgery, for example at the time of induction of anaesthesia.

#### **Cattle:**

Single subcutaneous or intravenous injection at a dosage of 0.5 mg meloxicam/kg body weight (i.e. 10 ml/100 kg body weight) in combination with antibiotic therapy or with oral re-hydration therapy, as appropriate.

#### **Pigs:**

Locomotor disorders:

Single intramuscular injection at a dosage of 0.4 mg meloxicam/kg body weight (i.e. 2 ml/25 kg body weight). If required, a second administration of meloxicam can be given after 24 hours.

Reduction of post-operative pain:

Single intramuscular injection at a dosage of 0.4 mg meloxicam/kg body weight (i.e. 0.4 ml/5 kg body weight) before surgery.

## **9. ADVICE ON CORRECT ADMINISTRATION**

Particular care should be taken with regard to the accuracy of dosing including the use of an appropriate dosing device and careful estimation of body weight.

Avoid introduction of contamination during use. The stopper should not be punctured more than 20 times.

## **10. WITHDRAWAL PERIOD(S)**

**Cattle:** meat and offal: 15 days

**Pigs:** meat and offal: 5 days

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special storage conditions.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after EXP.

Shelf-life after first opening the container: 28 days.

## **12. SPECIAL WARNING(S)**

### Special warnings for each target species:

Treatment of piglets with Meloxidolor before castration reduces post-operative pain. To obtain pain relief during surgery co-medication with an appropriate anaesthetic/sedative is needed.

To obtain pain relief for cattle and pigs during surgery co-medication with an appropriate anaesthetic/sedative/analgesic is needed.

To obtain the best possible pain relieving effect post-surgery Meloxidolor should be administered 30 minutes before surgical intervention.

Treatment of calves with Meloxidolor 20 minutes before dehorning reduces post-operative pain. Meloxidolor alone will not provide adequate pain relief during the dehorning procedure.

### Special precautions for use in animals:

If adverse reactions occur, treatment should be discontinued and the advice of a veterinarian should be sought.

Avoid use in any dehydrated, hypovolaemic or hypotensive animals, as there may be a potential risk of renal toxicity.

During anaesthesia, monitoring and fluid therapy should be considered as standard practice.

Any oral follow-up therapy using meloxicam or other NSAIDs should not be administered in cats, as appropriate dosage regimens for such follow-up treatments have not been established.

### Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Accidental self-injection may give rise to pain. People with known hypersensitivity to NSAIDs should avoid contact with the veterinary medicinal product.

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

The veterinary medicinal product should not be administered by pregnant women or women of child-bearing potential as Meloxicam may be harmful to the foetus and unborn child.

### Pregnancy and lactation:

**Dogs and cats:** Do not use in pregnant or lactating dogs or cats.

**Cattle:** Can be used during pregnancy.

**Pigs:** Can be used during pregnancy and lactation.

### Interactions with other medicinal products and other forms of interaction:

#### **For dogs and cats:**

Other NSAIDs, diuretics, anticoagulants, aminoglycoside antibiotics and substances with high protein binding may compete for binding and thus lead to toxic effects. Meloxidolor must not be administered in conjunction with other NSAIDs or glucocorticosteroids. Concurrent administration of potential nephrotoxic drugs should be avoided. In animals at anaesthetic risk (e.g. aged animals) intravenous or subcutaneous fluid therapy during anaesthesia should be taken into consideration. When anaesthesia and NSAID are concomitantly administered, a risk for renal function cannot be excluded.

Pre-treatment with anti-inflammatory substances may result in additional or increased adverse effects

and accordingly a treatment-free period with such veterinary medicinal products should be observed for at least 24 hours before commencement of treatment. The treatment-free period, however, should take into account the pharmacological properties of the products used previously.

**For cattle and pigs:**

Do not administer concurrently with glucocorticosteroids, other non-steroidal anti-inflammatory drugs or with anticoagulant agents.

Overdose (symptoms, emergency procedures, antidotes):

In case of overdose symptomatic treatment should be initiated.

Incompatibilities:

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

**13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

Detailed information on this veterinary medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu/>.

**15. OTHER INFORMATION**

**Package size:**

Colourless type I glass vial of 10 ml, 20 ml or 100 ml, closed with a rubber stopper and sealed with an aluminium cap.

Multi-packs of 5 x 20 ml and 10 x 20 ml.

Not all pack sizes may be marketed.

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

**België/Belgique/Belgien**

Kela Vetarinaria NV/SA  
Industriepark West 68  
B 9100 Sint-Niklaas  
Tel. : +32 3 7806390

**Luxembourg/Luxemburg**

Kela Vetarinaria NV/SA  
Industriepark West 68  
B 9100 Sint-Niklaas  
Tel. : +32 3 7806390

**Република България**

Vetpro Komers Ltd  
JK Trakia, bl 189, ent v, ap 2  
4000 Plovdiv  
Tel. +359 (0) 897 843918

**Magyarország**

Dopharma  
Zalmweg 24  
4941 VX Raamsdonksveer  
Tel. +31 162 582000

**Česká republika**

Sevaron ltd  
Palackeho trida 163 a  
61200 Brno  
Tel. +420 541 426 370

**Danmark**

Huvepharma NV  
Uitbreidingstraat 80  
B 2600 Antwerp - Belgium

**Deutschland**

Le Vet B.V.  
Wilgenweg 7  
NL 3421 TV Oudewater  
Tel. +31 348 565858

**Eesti**

AS Dimedium  
Emajõe 1a  
51008 Tartu  
Tel. +372 739 0660

**Ελλάδα**

Petline S.A.  
Finos Film Road, Thesi Xousmiza  
19004 Spata Attikis  
Tel. +030 210 6069800

**España**

Fatro Iberica  
C/Constitucion 1.BP.3  
8960 Sant Just Desvern  
Barcelona  
Tel: +34 934 802277

**France**

Osalia  
Rue Raffet 27-29  
75016 Paris  
Tel: +33 (0) 1 40505410

**Ireland**

NutriScience Ltd  
Unit 622  
Northern Extension IDA Industrial Estate  
Waterford  
Tel: +353 51 304010

**Malta**

Le Vet B.V.  
Wilgenweg 7  
NL 3421 TV Oudewater  
Tel. +31 348 565858

**Nederland**

Le Vet B.V.  
Wilgenweg 7  
NL 3421 TV Oudewater  
Tel. +31 348 565858

**Norge**

Le Vet B.V.  
Wilgenweg 7  
NL 3421 TV Oudewater  
Tel. +31 348 565858

**Österreich**

Vana GmbH  
Wolfgang Schmalzl-Gasse 6  
A 1020 Wien  
Tel. +43 1 728 03670

**Polska**

Drwalewskie Zaklady Przemyslu  
Bioweterynaryjnego S.A.  
Ul. Grojecka 6  
05-651 Drwalew  
Tel: +48 48 6649800

**Portugal**

Fatro Iberica  
C/Constitucion 1.BP.3  
8960 Sant Just Desvern  
Barcelona  
Tel: +34 934 802277

**România**

Dopharma  
Zalmweg 24  
4941 VX Raamsdonksveer  
Tel. +31 162 582000

**Slovenija**

Genera d.d  
Svetonedeljska 2 Kalinovica  
HR 10436 Rakov Potok  
Tel. +385 1 3388602

**Ísland**

Icevet  
Krokhalsi 14  
110 Reykjavik  
Sími: +354-820 2240

**Italia**

PH Farmaceutici  
Via Mazzini 3/C  
20063 Cernusco Sul Naviglio, Milano  
Tel: +39 (0)2 2222 3781

**Κύπρος**

Petline S.A.  
Finos Film Road, Thesi Xousmiza  
19004 Spata Attikis  
Tel. +030 210 6069800

**Latvija**

AS Dimedium  
Emajõe 1a  
51008 Tartu  
Tel. +372 739 0660

**Lietuva**

AS Dimedium  
Emajõe 1a  
51008 Tartu  
Tel. +372 739 0660

**Slovenská republika**

Sevaron ltd  
Palackeho trida 163 a  
61200 Brno  
Tel. +420 541 426 370

**Suomi/Finland**

FaunaPharma  
Orionintie 5  
Pääovi, 3. krs.  
02200 Espoo  
Puh/Tel: +358 45 2300665

**Sverige**

VM Pharma AB  
Postbox 45010  
SE-140 30 Stockholm  
info@vetmedic.se

**United Kingdom**

NutriScience Ltd  
Unit 622  
Northern Extension IDA Industrial Estate  
Waterford  
Tel: +353 51 304010

**Republika Hrvatska**

Genera d.d  
Svetonedeljska 2 Kalinovica  
HR 10436 Rakov Potok  
Tel. +385 1 3388602



## PACKAGE LEAFLET FOR:

### Meloxidolor 20 mg/ml solution for injection for cattle, pigs and horses

#### **1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder:

Le Vet Beheer B.V.  
Wilgenweg 7  
3421 TV Oudewater  
The Netherlands

Manufacturer for the batch release:

Produlab Pharma B.V.  
Forellenweg 16  
4941 SJ Raamsdonksveer  
The Netherlands

#### **2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Meloxidolor 20 mg/ml solution for injection for cattle, pigs and horses  
meloxicam

#### **3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENTS**

One ml contains:

**Active substance:**

Meloxicam 20 mg

**Excipient:**

Ethanol 150 mg

Clear yellow solution.

#### **4. INDICATIONS**

**Cattle:**

For use in acute respiratory infection with appropriate antibiotic therapy to reduce clinical signs in cattle.

For use in diarrhoea in combination with oral re-hydration therapy to reduce clinical signs in calves of over one week of age and young, non-lactating cattle.

For adjunctive therapy in the treatment of acute mastitis, in combination with antibiotic therapy.

For the relief of post-operative pain following dehorning in calves.

**Pigs:**

For use in non-infectious locomotor disorders to reduce the symptoms of lameness and inflammation.

For adjunctive therapy in the treatment of puerperal septicaemia and toxemia (mastitis-metritis-agalactia syndrome) with appropriate antibiotic therapy.

**Horses:**

For use in the alleviation of inflammation and relief of pain in both acute and chronic musculo-skeletal disorders.

For the relief of pain associated with equine colic.

**5. CONTRAINDICATIONS**

Do not use in horses less than 6 weeks of age.

Do not use in animals suffering from impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of ulcerogenic gastrointestinal lesions.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

For the treatment of diarrhoea in cattle, do not use in animals of less than one week of age.

See also Section 12.

**6. ADVERSE REACTIONS**

In cattle and pigs, subcutaneous, intramuscular as well as intravenous administration is well tolerated; only a slight transient swelling at the injection site following subcutaneous administration was observed in less than 10 % of the cattle treated in clinical studies.

In horses, in rare cases a transient swelling at the injection site can occur but resolves without intervention.

In very rare cases anaphylactoid reactions, which may be serious (including fatal), may occur and should be treated symptomatically.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reactions)
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

**7. TARGET SPECIES**

Cattle, pigs and horses

**8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION****Cattle:**

Single subcutaneous or intravenous injection at a dosage of 0.5 mg meloxicam/kg body weight (i.e. 2.5 ml/100 kg body weight) in combination with antibiotic therapy or with oral re-hydration therapy, as appropriate.

**Pigs:**

Single intramuscular injection at a dosage of 0.4 mg meloxicam/kg body weight (i.e. 2 ml/100 kg body weight) in combination with antibiotic therapy, as appropriate. If required, a second administration of meloxicam can be given after 24 hours.

**Horses:**

Single intravenous injection at a dosage of 0.6 mg meloxicam/kg body weight (i.e. 3 ml/100 kg body weight).

Avoid introduction of contamination during use. The stopper should not be punctured more than 20 times.

**9. ADVICE ON CORRECT ADMINISTRATION**

Avoid introduction of contamination during use. The stopper should not be punctured more than 20 times.

**10. WITHDRAWAL PERIOD(S)**

**Cattle:** meat and offal: 15 days; milk: 5 days

**Pigs:** meat and offal: 5 days

**Horses:** meat and offal: 5 days.

Not authorised for use in horses producing milk for human consumption.

**11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special storage conditions.

Do not use after the expiry date stated on the label after EXP.

Shelf-life after first opening the container: 28 days.

**12. SPECIAL WARNING(S)**

Special warnings for each target species:

Treatment of calves with Meloxidolor 20 minutes before dehorning reduces post-operative pain. Meloxidolor alone will not provide adequate pain relief during the dehorning procedure. To obtain adequate pain relief during surgery co-medication with an appropriate analgesic is needed.

Special precautions for use in animals:

If adverse reactions occur, treatment should be discontinued and the advice of a veterinarian should be sought.

Avoid use in very severely dehydrated, hypovolaemic or hypotensive animals which require parenteral rehydration, as there may be a potential risk of renal toxicity.

In case of inadequate relief of pain when used in the treatment of equine colic, careful re-evaluation of the diagnosis should be made as this could indicate the need for surgical intervention.

Special precautions to be taken by the person administering the veterinary medicinal product to

animals: Accidental self-injection may give rise to pain. People with known hypersensitivity to Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) should avoid contact with the veterinary medicinal product.

In case of accidental self-injection, seek medical advice immediately and show this package leaflet or the label to the physician.

The veterinary medicinal product should not be administered by pregnant women or women of child-bearing potential as Meloxicam may be harmful to the foetus and unborn child.

Pregnancy and lactation:

**Cattle and pigs:** Can be used during pregnancy and lactation.

**Horses:** Do not use in pregnant or lactating mares.

Interactions with other medicinal products and other forms of interaction:

Do not administer concurrently with glucocorticosteroids, other non-steroidal anti-inflammatory drugs or with anticoagulant agents.

Overdose (symptoms, emergency procedures, antidotes)

In case of overdose, symptomatic treatment should be initiated.

Incompatibilities:

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

**13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

Detailed information on this veterinary medicinal product is available on the website of the European Medicines Agency (<http://www.ema.europa.eu>).

**15. OTHER INFORMATION**

**Package (size)**

Colourless type I glass vials of 50 ml or 100 ml, closed with a rubber stopper and sealed with an aluminium cap.

Multi-pack of 12 x 100ml

Not all pack sizes may be marketed.

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

**België/Belgique/Belgien**

Kela Vetarinaria NV/SA  
Industriepark West 68  
B 9100 Sint-Niklaas  
Tel. : +32 3 7806390

**Luxembourg/Luxemburg**

Kela Vetarinaria NV/SA  
Industriepark West 68  
B 9100 Sint-Niklaas  
Tel. : +32 3 7806390

**Република България**

Vetpro Komers Ltd  
JK Trakia, bl 189, ent v, ap 2  
4000 Plovdiv  
Tel. +359 (0) 897 843918

**Magyarország**

Dopharma  
Zalmweg 24  
4941 VX Raamsdonksveer  
Tel. +31 162 582000

**Česká republika**

Sevaron ltd  
Palackeho trida 163 a  
61200 Brno  
Tel. +420 541 426 370

**Danmark**

Huvepharma NV  
Uitbreidingstraat 80  
B 2600 Antwerp - Belgium

**Deutschland**

Le Vet B.V.  
Wilgenweg 7  
NL 3421 TV Oudewater  
Tel. +31 348 565858

**Eesti**

AS Dimedium  
Emajõe 1a  
51008 Tartu  
Tel. +372 739 0660

**Ελλάδα**

Petline S.A.  
Finos Film Road, Thesi Xousmiza  
19004 Spata Attikis  
Tel. +030 210 6069800

**España**

Fatro Iberica  
C/Constitucion 1.BP.3  
8960 Sant Just Desvern  
Barcelona  
Tel: +34 934 802277

**France**

Osalia  
Rue Raffet 27-29  
75016 Paris  
Tel: +33 (0) 1 40505410

**Ireland**

NutriScience Ltd  
Unit 622  
Northern Extension IDA Industrial Estate  
Waterford  
Tel: +353 51 304010

**Malta**

Le Vet B.V.  
Wilgenweg 7  
NL 3421 TV Oudewater  
Tel. +31 348 565858

**Nederland**

Le Vet B.V.  
Wilgenweg 7  
NL 3421 TV Oudewater  
Tel. +31 348 565858

**Norge**

Le Vet B.V.  
Wilgenweg 7  
NL 3421 TV Oudewater  
Tel. +31 348 565858

**Österreich**

Vana GmbH  
Wolfgang Schmalzl-Gasse 6  
A 1020 Wien  
Tel. +43 1 728 03670

**Polska**

Drwalewskie Zaklady Przemyslu  
Bioweterynaryjnego S.A.  
Ul. Grojecka 6  
05-651 Drwalew  
Tel: +48 48 6649800

**Portugal**

Fatro Iberica  
C/Constitucion 1.BP.3  
8960 Sant Just Desvern  
Barcelona  
Tel: +34 934 802277

**România**

Dopharma  
Zalmweg 24  
4941 VX Raamsdonksveer  
Tel. +31 162 582000

**Slovenija**

Genera d.d  
Svetonedeljska 2 Kalinovica  
HR 10436 Rakov Potok  
Tel. +385 1 3388602

**Ísland**

Icevet  
Krokhalsi 14  
110 Reykjavik  
Sími: +354-820 2240

**Italia**

PH Farmaceutici  
Via Mazzini 3/C  
20063 Cernusco Sul Naviglio, Milano  
Tel: +39 (0)2 2222 3781

**Κύπρος**

Petline S.A.  
Finos Film Road, Thesi Xousmiza  
19004 Spata Attikis  
Tel. +030 210 6069800

**Latvija**

AS Dimedium  
Emajõe 1a  
51008 Tartu  
Tel. +372 739 0660

**Lietuva**

AS Dimedium  
Emajõe 1a  
51008 Tartu  
Tel. +372 739 0660

**Slovenská republika**

Sevaron ltd  
Palackeho trida 163 a  
61200 Brno  
Tel. +420 541 426 370

**Suomi/Finland**

FaunaPharma  
Orionintie 5  
Pääovi, 3. krs.  
02200 Espoo  
Puh/Tel: +358 45 2300665

**Sverige**

VM Pharma AB  
Postbox 45010  
SE-140 30 Stockholm  
info@vetmedic.se

**United Kingdom**

NutriScience Ltd  
Unit 622  
Northern Extension IDA Industrial Estate  
Waterford  
Tel: +353 51 304010

**Republika Hrvatska**

Genera d.d  
Svetonedeljska 2 Kalinovica  
HR 10436 Rakov Potok  
Tel. +385 1 3388602